
COMPASS[®] ENTEROBACTER SAKAZAKII AGAR

DETECTION OF *CRONOBACTER SAKAZAKII*

1 INTENDED USE

The COMPASS[®] *Enterobacter sakazakii* Agar is used for the detection of *Cronobacter sakazakii* and spp. in milk powder, dehydrated products and their components found in infant foods.

The type composition of the Chromogenic *Cronobacter* Isolation Agar conforms to the formulation found in the project directive PR NF EN ISO 22964.

2 HISTORY

Cronobacter sakazakii (formerly *Enterobacter sakazakii*) is a Gram negative bacillus, mobile, non-sporulated facultative anaerobe which forms pigmented yellow colonies after 48-72 hours of incubation on non-selective media. An opportunistic pathogen, it is notably at the origin of meningitis and enteritis, particularly with newborns and young children, and although the frequency is rather low at 1 in 100000, the mortality is high at roughly 20 to 50%. While the strains have been isolated from different food products, only those products destined for infant or baby foods are implicated in the infectious episodes.

Studies have shown that 100% of the *Cronobacter sakazakii* were positive for α -glucosidase when at the same time 100% of other species of *Enterobacter* were negative for this enzyme. On the basis of these observations, the chromogenic substrate 5-bromo-4-chloro-3-indolyl- α -D-glucopyranoside (X- α -glucoside) has been proposed for differentiating this strains from other members of the *Enterobacteriaceae* family.

3 PRINCIPLES

Tryptone stimulates the growth of *Cronobacter*.

Yeast extract is a source of complex vitamin B.

Sodium chloride maintains osmotic pressure.

The choice of the incubation temperature fixed at 44 °C, plus the association of sodium desoxycholate and crystal violet combines to inhibit the growth of a large spectrum of contaminating microflora.

The enzyme α -glucosidase hydrolyzes the X- α -glucoside and liberates the aglycone 5 bromo-4-chloro-indolol. In the presence of oxygen, this aglycone is dimerized and forms the pigment bromo-chloro-indigo.

4 TYPICAL COMPOSITION

The composition can be adjusted in order to achieve optimal performance.

For 1 liter of media :

- Tryptone	7,00 g
- Yeast extract	3,00 g
- Sodium chloride	5,00 g
- Sodium desoxycholate	0,60 g
- Cristal violet.....	2,0 mg
- 5-bromo-4-chloro-3-indolyl, α -D-glucopyranoside	150,0 mg
- Bacteriological agar.....	14,40 g

pH of the ready-to-use media at 25 °C : 7,0 \pm 0,2.

5 PREPARATION

- Dissolve 30,2 g of dehydrated media (BK188) in 1 liter of distilled or demineralized water.
- Slowly stir until complete dissolution.
- Divide into vials.
- Sterilize in an autoclave at 121 °C for 15 minutes.
- Cool to 44-47 °C.
- Pour into sterile Petri plates and let solidify on a cold surface.

✓ **Reconstitution :**
30,2 g/L

✓ **Sterilization :**
15 min at 121 °C

6 INSTRUCTIONS FOR USE

- Dry the plates in an incubator with the covers partially removed.
- On the surface of plates prepared as above, or on pre-poured plates (BM120) brought to room temperature, inoculate by streaking a loop of enrichment media (mLST broth, BM121).
- Incubate at 44 ± 1°C for 24 ± 2 hours.

✓ **Inoculation :**
a loop of enrichment media

✓ **Incubation :**
24 h at 44°C

7 RESULTS

The aspect of the colonies are as follows :

Microorganisms	Colony characteristics
<i>Cronobacter sakazakii</i>	Blue-violet colonies
<i>Escherichia coli</i>	Grey to violet colonies
<i>Enterobacter</i> spp., <i>Klebsiella</i> spp.	Grey to violet colonies
Gram positive bacteria	Inhibited

See ANNEX 1 : PHOTO SUPPORT.

Note : The strains of *Cronobacter* present generally a characteristic pigmentation blue-green. However, it is possible that some strains weakly α -glucosidase-positives could be confused with certain non-targeted Gram-negative microorganisms, which by incorporating the crystal violet, present a similar aspect.

8 QUALITY CONTROL

Dehydrated media : beige powder, free-flowing and homogeneous.

Prepared media : violet agar.

Typical culture response after 24 of incubation at 44 °C:

Microorganisms	Growth	Characteristics
<i>Cronobacter sakazakii</i> WDCM 00214	Good, score 2	Blue-green colonies
<i>Cronobacter sakazakii</i> CIP 104951	Good, score 2	Blue-green colonies
<i>Escherichia coli</i> WDCM 00013	Good, score 2	Violet colonies
<i>Enterobacter cloacae</i> WDCM 00083	Good, score 2	Violet colonies
<i>Staphylococcus aureus</i> WDCM 00034	Inhibited, score 0	-

9 STORAGE / SHELF LIFE

Dehydrated base media : 2-30 °C.

Pre-poured media in Petri plates : 2-8 °C.

The expiration date is indicated on the label.

Prepared media in Petri plates (*) : 30 days at 2-8 °C.

(*) Benchmark value, determined in standard conditions of preparation, following manufacturer's instructions.

10 PACKAGING

Dehydrated media :

500 g bottle BK188HA

Pre-poured media in Petri plates (Ø 90 mm) :

20 plates BM12008

11 BIBLIOGRAPHY

Muytjens, H.L., van der ROS, van de Repe, J., and van Druuten, H.A.. 1984. Enzymatic profiles of *Enterobacter sakazakii* and related species with special reference to the alpha-glucosidase reaction and reproducibility of the test system. *Journal of Clinical Microbiology*, 20 : 684-686.

Simmons, B.P., Gelfand, M.S., Haas, M., Metts, L., and Ferguson, J.. 1989. *Enterobacter sakazakii* infections in neonates associated with intrinsic contamination of a powdered infant formula. *Infection Control and Hospital Epidemiology*, 10 : 398-401.

Iversen, C., Drugan, P., and Forsythe, S.. 2004. A selective differential medium for *Enterobacter sakazakii*, a preliminary study. *International Journal of Food Microbiology*, 96 : 133-139.

Lehner, A., and Stephan, R.. 2004. Microbiological, epidemiological and food safety aspects of *Enterobacter sakazakii*. *Journal of Food Protection*, 67(12) : 2850-2857.

Guillaume-Gentil, O., Sànnard, V., Kandhai, M.C., Marugg, J.D., and Joosten, H.. 2005. A simple and rapid cultural method for detection of *Enterobacter sakazakii* in environmental samples. *Journal of Food Protection*, 68(1) : 64-69.

Gurtler, J.B., Kornacki, J.L., and Beuchat, L.R.. 2005. *Enterobacter sakazakii*: a coliform of increased concern to infant health. *International Journal of Food Microbiology*, 104 : 1-34.

ISO/TS 22964. Février 2006. Lait et produits laitiers. Détection de l'*Enterobacter sakazakii*.

12 ADDITIONAL INFORMATION

The information provided on the labels take precedence over the formulations or instructions described in this document and are susceptible to modification at any time, without warning.

Document code : COMPASS ENTEROBACTER SAKAZAKII_ENv4

Creation date : 02-2006

Updated : 03-2016

Origin of revision : General update.


ANNEX 1 : PHOTO SUPPORT

COMPASS[®] *Enterobacter sakazakii* Agar

Detection of *Cronobacter sakazakii*.

Results :

Growth obtained after 24 hours of incubation at 44 °C.



Cronobacter sakazakii

Characteristic colony :
blue-green color